



SOCIEDAD ESPAÑOLA DE
CALIDAD ASISTENCIAL

Revista de Calidad Asistencial

www.elsevier.es/calasis



ORIGINAL ARTICLE

Improving quality in healthcare: What makes a satisfied patient?



A. Más^{a,*}, P. Parra^a, R.M. Bermejo^b, M.D. Hidalgo^b, J.E. Calle^a

^a Programa EMCA, Subdirección General de Planificación, Calidad e Investigación, Consejería de Sanidad, Región de Murcia, Spain

^b Universidad de Murcia, Murcia, Spain

Received 10 September 2015; accepted 26 November 2015

Available online 8 February 2016

KEYWORDS

Patient satisfaction;
Quality of care;
Satisfaction in
healthcare;
Validity;
Survey;
Questionnaires

Abstract

Objective: To update the metric properties of a perceived quality questionnaire for patients achieved, and to identify the variables which predict satisfaction.

Methods: Self-administered questionnaire completed at home following patient discharge, using a questionnaire prepared by the authors on a sample of 7207 users of medical departments in 9 public hospitals during the years 2006–2009. A principal component analysis with varimax rotation was performed. Reliability was assessed using internal consistency coefficient. An analysis was made of the compliance with each indicator reported by respondents. A logistic regression analysis was performed to determine the perceived quality dimensions which predicted overall patient satisfaction.

Results: The results of the reliability analysis indicated good coefficients for interpersonal manner (0.94) and professional competence (0.85) dimensions, and moderate values for the other dimensions (comfort 0.55, information 0.38, and organisation 0.37). Factor analyses showed single factors in each of the perceived quality dimensions, with a percentage of explained variance greater than 35% for information, interpersonal manner, professional competence, and comfort, and less than 30% for organisation. The dimensions which predicted satisfaction were interpersonal manner of healthcare staff, professional competence, and information.

Conclusions: The metric properties of the questionnaire used have been updated, yielding a valid and reliable questionnaire for assessing patient satisfaction in quality management programmes, both for internal purposes and for conducting external comparisons.

A positive relationship was obtained between the level of patient satisfaction and level of professional competence, interpersonal manner of healthcare staff, and information received.

© 2016 SECA. Published by Elsevier España, S.L.U. All rights reserved.

* Corresponding author.

E-mail address: adelia.mas@carm.es (A. Más).

PALABRAS CLAVE

Satisfacción del paciente;
Calidad de atención;
Satisfacción en la asistencia sanitaria;
Validez;
Encuesta;
Cuestionarios

Mejora de la calidad en los cuidados de salud: ¿Qué hace que un paciente esté satisfecho?**Resumen**

Objetivo: Revisar las propiedades métricas de un cuestionario de calidad percibida de los servicios médicos hospitalarios, determinar el nivel de satisfacción de los usuarios e identificar los indicadores de calidad predictores de satisfacción.

Métodos: Se ha utilizado un cuestionario autoadministrado en el domicilio al alta del paciente. La muestra fue de 7.207 usuarios de servicios médicos en 9 hospitales públicos desde 2006 a 2009. Se analizó la fiabilidad (consistencia interna) de las dimensiones de calidad percibida y su validez interna usando análisis de componentes principales. Finalmente, se realizó un análisis de regresión logística para determinar las dimensiones de calidad que mejor predicen la satisfacción del paciente.

Resultados: Las dimensiones de trato y competencia profesional mostraron altos coeficientes de fiabilidad (0,94 y 0,85, respectivamente), para el resto de dimensiones los valores fueron moderados (0,37 a 0,55). El análisis de componentes principales indicó que las diferentes dimensiones de calidad (trato, competencia profesional, información, confort y organización) son unidimensionales, explicando cada una de ellas más de un 35% de la varianza, excepto para organización, que fue próxima al 30%. El trato recibido por el usuario, la competencia profesional percibida y la información fueron las dimensiones de calidad que mejor pronosticaron la satisfacción del paciente.

Conclusiones: El cuestionario de calidad percibida mostró adecuadas propiedades métricas, siendo válido y fiable para la valoración de la satisfacción del paciente en los programas de gestión de la calidad, tanto con fines internos como para la realización de comparaciones externas.

© 2016 SECA. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

Introduction

Information on the degree of satisfaction and perceived quality reported by health service users and their families is useful for improving the organisation of healthcare, and serves as a starting point for improving service quality.¹⁻⁴ Patient satisfaction forms part of health centre outcomes and is an essential element in the evaluation of these organisations. Hence, the dimension of patient satisfaction with care received is included in all healthcare quality assessment models.

However, patient satisfaction is a complex variable⁴ and there is no general consensus on the factors with which it is associated. In a systematic review⁵ of over 3000 articles, the authors concluded that satisfaction is a relative concept, a reflection of patient assessments of the quality of care they received compared to subjective standards, which can be influenced by previous health status, severity of illness or other factors such as age. They also indicated inconsistent influence of gender, race, socioeconomic or educational level on patient satisfaction. Furthermore, satisfaction is a multidimensional concept which varies according to the type of care provided and which requires the use of different tools for the different areas of health care.⁶

Of the various methods available to determine the degree of satisfaction with health services, questionnaires are most commonly used.⁵ Among their advantages, questionnaires can be administered to any group of people, information about past events can be retrieved, a large amount of

information can be collected in a short time and they are relatively inexpensive.⁷

However, most of the questionnaires used to measure perceived quality and patient satisfaction did not provide data about metric characteristics (reliability and validity).⁸ In others it is necessary to update their metric properties, this is a necessary procedure since the characteristics of the target population may change over time. In addition, many of the questionnaires used to measure both perceived quality and inpatient satisfaction are the same for all patients, regardless of the illness that has resulted in their hospital admission or the unit in which they are cared for. Thus, patients admitted in medical services present characteristics that differentiate them from patients admitted to other hospital units,^{9,10} since they are older,¹¹ usually present comorbidity¹²⁻¹⁵ and have a longer average stay and a higher possibility of readmission.

The aim of this study was to update the metric properties of a perceived quality questionnaire used in the Hospital Medical Services,¹⁶ determine the degree of patient satisfaction with the care received and the variables related with patient satisfaction.

Material and methods**Participants**

The study population comprised the total number of patients discharged from the medical service of the nine general

hospitals run by the Murcian Health Service (MHS) over the period 2006–2009, from which a sample of 7207 cases was extracted. The MHS is the public body responsible for the provision of health services in the Autonomous Region of Murcia that, located in the south-east Spain had, at the time of the study, a reference population of 1,446,109 inhabitants and nine general hospitals. The size of these nine hospitals were from 97 to 863 beds. Of the nine, one had more than 500 beds (large hospital), 4 had a medium size (200–500 beds) and the other four had less than 200 (small).

From 2006 to 2009, data was collected four times a year, once every three months. In order to identify cases, a period of between one and three weeks each quarter, depending on the size of the hospital, was selected. A stratified random start systematic sampling, not proportional to hospital size, was conducted.

Questionnaire

We used the EMCA (*Assessing and Improving Health-care Quality*) questionnaire to measure satisfaction and perceived quality during patient hospitalisation in medical services. This questionnaire was developed following the methodology used by the Picker Institute (Boston, USA),¹⁷ which includes an exploration with focus groups in order to identify those relevant aspects to patients admitted to hospital.

The statements were formulated in accordance with recommendations for questionnaire design.¹⁸ Subsequently, a pilot study and preliminary validation was performed using patients for different regions in Spain.

Three different questionnaires were developed: one for patients admitted for surgery services, another for women admitted to give birth, and the last, which was the focus of this research, a specific questionnaire to explore the opinions of patients admitted to Hospital Medical Services.

The questionnaire for measuring perceived quality and satisfaction of patients admitted to medical services included five perceived quality dimensions: interpersonal manner, information, professional competence, organisation and comfort. It consisted of fifty-one items on perceived quality, one item to determine overall assessment of satisfaction using a 10-point response scale from 0 to 10 (from very dissatisfied to very satisfied) and four items referring to patient profile (age, gender, educational level and previous hospital admissions).¹⁶

To facilitate understanding and self-completion of the tool, the order of the questions followed the sequence of the care process itself (admission, length of stay and discharge).

Data collection

The patient received the questionnaire at home, accompanied by a cover letter, within ten days of discharge. Patients returned the completed questionnaire using a postage-paid envelope. To increase the response rate, two reminder letters were sent 10 and 20 days after the questionnaire had initially been posted. The response rate was 34.92%.

Data analyses

A total of 2517 questionnaires were received. Of this total 1789 (71%) were excluded from the psychometric analysis because their items answers were incomplete. The possible existence of sample selection bias was assessed by comparing the mean age and gender final sample distribution with respect to the initial sample. Given the large sample size, the magnitude of the mean difference using Cohen's *d* test¹⁹ was calculated. Furthermore, univariate descriptive analyses of the sociodemographic variables were conducted.

In order to assess dimensional structure principal component analysis with varimax rotation was carried out. This analysis provided information on internal questionnaire validity. Reliability was calculated using Chronbach's internal consistency coefficient.

Then we calculated the percentage of compliance with the perceived quality variables by dividing the number of positive responses for each variable by the total number of responses received for each. An indicator was considered satisfactory according to whether the respondent answered "yes" or "no", or in the case of multiple choice items, indicated the extreme value of the statements, namely "always", "never" or "very good". Also composite measures for each of the dimensions obtained in the factor analysis, using the opportunity model, as the sum of positive responses to the items in each dimension divided by the sum of the answers to those same items²⁰ were calculated.

Lastly, a logistic regression analysis in order to determine the perceived quality dimensions that best predicted overall patient satisfaction was performed. Since our aim was to determine the level of quality perceived by very satisfied users, patients were classified into two categories: patients who reported a very high level of satisfaction (score of 9 or above for the overall satisfaction criterion) and other patients (score of 8 or below). In order to control for confounding factors, the model also included sociodemographic variables, previous hospital admissions and hospital size (large, medium and small).

Statistical analysis of the data was performed using SPSS 19.0.²¹ In all cases, results were considered statistically significant when the probability value was less than or equal to 0.05.

Results

The mean age of study participants was 70.2 years old, 45.2% were female and 84.1% didn't had studies or had completed primary education (Table 1).

Comparing the sociodemographic characteristic of analysed sample with the initial sample, no significant differences were found for gender distribution. However, statistical significant differences were found for age ($p < 0.001$), although the effect size was small (Cohen's $d < 0.20$) indicating that the differences were negligible.

Dimensional structure analysis

Using the Kaiser–Meyer–Olkin (KMO) index of sampling adequacy, a value of 0.89 for Information dimension, 0.93 for Interpersonal Manner, 0.86 for professional competence,

Table 1 Profile of the study sample.

	<i>n</i>	%
Sex		
Male	399	54.8
Female	329	45.2
Age		
18–44 (1)	58	8.0
45–64 (2)	144	19.8
>64 (3)	526	72.3
Educational level		
No formal education/primary education (1)	582	84.1
Secondary education/vocational training (2)	92	13.3
Diploma/degree (3)	18	2.6

0.59 for organisational dimension and 0.70 for comfort were obtained, indicating that the data were suitable for performing factor analyses. Bartlett's sphericity test results were statistically significant for each of the dimensions ($p < 0.001$).

The factor analysis results for the 12 items related to Information showed that a single factor explained 36.35% of the total variance (eigenvalue = 4.36). This dimension included statements that referred to the information received at the time of admission, during stay in the emergency department and on arrival at the ward, easily understood answers to questions directed at medical or nursing staff, explanations of the patient's condition during his or her stay on the ward given to the patient or his or her companions and discharge information. The 17 items related to interpersonal dimension, were grouped into a single factor explained 39.79% of the variance (eigenvalue = 6.77). This dimension comprised statements that referred to issues related to respect for privacy, absence of discriminatory treatment and kindness. Moreover, the 8 items related to professional competence were grouped into a single factor explained 50.68% of the total variance (eigenvalue = 4.05). This dimension included statements referring to the professionalism of all categories of hospital staff, absence of care errors and approaches to pain relief. The five items for the dimension of organisational, which referred to compliance with rules, shifts and visits, revealed a single factor structure explaining 29.14% of the total variance (eigenvalue = 1.46). Lastly, the five items related to comfort were

grouped into a single factor which explained 36.13% of the total variance (eigenvalue = 1.81). The items in this dimension referred to the condition of the premises and the comfort of the hospital's medical services.

The minimum and maximum factor loadings for each of the dimensions are shown in Table 2.

Reliability of the test

The internal consistency coefficient established by Cronbach's alpha coefficient³¹ for each of the dimensions was 0.83 (information), 0.90 (interpersonal manner), 0.85 (professional competence), 0.37 (organisational) and 0.55 (comfort).

Cronbach's alpha coefficient for all questionnaire items was 0.94.

Percentage of compliance with perceived quality items

The percentages of compliance obtained for each dimension ranged from 74.3% for the dimension of organisational to 59.5% for the professional competence dimension (see Table 3). Percentages for each item ranged between 93.4% (clear information about the reason for admission) to 42.3% (professionalism of cleaning staff).

Overall satisfaction rating and logistic regression analysis

Mean overall satisfaction on a scale of 0–10 was 7.99, with 43.6% of patients giving a score of 9–10. In order to determine the variables that best predicted the degree of satisfaction of hospitalised patients, a logistic regression analysis was performed, using patient satisfaction as the dependent variable and the perceived quality dimensions (information, interpersonal manner, professional competence, organisational and comfort), socio-demographic variables (educational level, age and gender of the patient), previous admissions to hospital and hospital size (large, medium and small) as independent variables.

The overall fit of this model was statistically significant (Chi-square = 324.39, $df = 10$, $p < 0.01$; Nagelkerke's $R^2 = 0.523$) and it correctly classified 80.5% of cases. Information, interpersonal manner and professional competence

Table 2 Range of factor loadings.

Dimension	Item with minimum factor loading value		Item with maximum factor loading value	
Information	Information about hospital rules	0.33	Clear explanation of the patient's progress	0.76
Interpersonal manner	Problems related to interpersonal manner	0.48	Interpersonal manner of nursing staff	0.75
Professional competence	Treatment, test or care errors	0.32	Professionalism of auxiliary staff and porters	0.89
Organisation	Medical care from the doctor	0.46	Problems with visits	0.65
Comfort	No lack of towels or bed linen	0.45	State of hospital units	0.68

Table 3 Percentage of satisfaction by item and dimension.

Item	Indicators	% Satisfaction
1	Information about hospital rules	69.0
5	Explanation in the emergency department about what they were doing	52.2
9	Sufficiency of information given to patient's companions in the emergency department	51.2
11	Clear information on the reason for admission	93.4
23	Clear explanation of the patient's progress	67.9
25	Possibility of discussing aspects of the illness with a doctor	74.7
27	Easily understood responses from doctors	69.4
28	Easily understood responses from nursing staff	64.4
29	Sufficient information given to patient's companions	68.1
30	No different or contradictory information given by staff	92.7
46	Clear explanation on discharge about treatment and care to observe once at home	89.4
47	Possibility of asking doctors questions about the discharge report	87.6
	<i>Dimension of information</i>	73.3
16	No problems related to room sharing	72.8
17	Quiet visits	82.0
38	Compliance with no smoking rules	77.9
41	Care not affected by shift changes	82.0
43	Medical care from the same doctor	56.6
	<i>Dimension of organisation</i>	74.3
6	Interpersonal manner helped patients to feel calmer and more relaxed	58.5
7	Gentle manner during physical examinations	75.4
8	Kind manner to patient's companions in the emergency department	64.3
12	Easy identification of staff	67.2
13	The staff responsible for care introduced themselves	45.5
14	No problems related to interpersonal manner	87.1
15	Help received for personal hygiene tasks	69.8
18	Availability of staff whenever needed	81.5
24	Patient's opinions taken into account	63.6
33	Respect for privacy during physical examinations, personal hygiene or wound dressing	86.3
34	Support from medical or nursing staff when experiencing anxiety or depression	63.0
35	Rapid attention to patient needs	52.9
45_1	Interpersonal manner of medical staff	84.2
45_2	Interpersonal manner of nursing staff	69.5
45_3	Interpersonal manner of nursing auxiliaries	68.0
45_4	Interpersonal manner of porters	69.6
45_5	Interpersonal manner of cleaning staff	73.4
	<i>Dimension of interpersonal manner</i>	69.4
26	No treatment, test or care errors	86.5
39	The work carried out by the cleaning staff was not bothersome	73.8
40	The necessary means were employed to relieve pain	74.2
51_1	Professionalism of medical staff	63.0
51_2	Professionalism of nursing staff	48.2
51_3	Professionalism of auxiliary staff	44.5
51_4	Professionalism of porters	43.1
51_5	Professionalism of cleaning staff	42.3
	<i>Dimension of professional competence</i>	59.5
10	Everything in the hospital emergency department was in good condition	58.0
19	No problems with food	54.0
20	No lack of towels or bed linen	90.4
21	Hospital units were in good condition	48.1
37	No problems in use of showers and toilets	67.4
	<i>Dimension of comfort</i>	63.6

Table 4 Results of logistic regression analysis.

Variable	$\leq 8^{a,b}$	$\geq 9^{a,b}$	β	Wald	<i>p</i>
Interperson	62.28 (25.22)	87.41 (16.83)	0.016	5.383	0.020
Interpersonal manner	55.08 (27.53)	87.56 (15.67)	0.036	22.765	0.000
Competence	42.90 (28.14)	80.75 (26.25)	0.025	37.198	0.000
Organisation	68.35 (24.51)	81.47 (19.46)	0.001	0.008	0.929
Comfort	55.16 (29.35)	74.38 (25.20)	0.006	2.177	0.140
Educational level	1 (55.9%)	1 (44.1%)	0.044	0.026	0.871
	2 (57.6%)	2 (42.4%)			
	3 (72.2%)	3 (27.8%)			
Age	1 (63.8%)	1 (36.2%)	0.169	0.817	0.366
	2 (51.1%)	2 (48.9%)			
	3 (57.0%)	3 (43.0%)			
Hospital size	Small (46.4%)	Small (53.6%)	-0.294	3.559	0.059
	Medium (57.2%)	Medium (42.8%)			
	Large (58.3%)	Large (41.7%)			
Sex	Male (55.2%)	Male (44.8%)	-0.276	1.649	0.199
	Female (57.8%)	Female (42.2%)			
Previous hospital admissions	Yes (48.9%)	Yes (51.1%)	0.147	0.372	0.542
	No (58.7%)	No (41.3%)			
Constant			-6.034	36.527	0.000

^a Means and standard deviations for the dimensions.

^b Percentages for socio-demographic and hospital variables.

variables were statistically significant, whilst Hospital size was partially significant (see Table 4).

The final model including these four significant variables correctly classified 80.4% of cases (Chi-square = 350.02, *df* = 4, *p* < 0.01).

Discussion

In this study, the metric properties of a questionnaire aimed at assessing patients' perceptions of the care received in hospital medical services were revised. Reliability and construct validity were analysed, obtaining updated data about dimensionality questionnaire. These perceived quality dimensions were one-dimensional.

Cronbach's alpha coefficients were very satisfactory for most scales. Reliability coefficients were high for the dimensions of information, interpersonal manner and professional competence, and thus the overall score obtained with this questionnaire will serve to conduct comparisons between different hospital medical services, which is one of the objectives of its use. The value obtained for the organisational dimension was low compared to that obtained in the study by Granado et al. for the factor measuring "Administrative Quality".²² This difference could be due to the fact that the response scale used in the study by Granado had a greater score range, as each item presented 10 response categories. The internal consistency of the comfort dimension was moderate. Thus, in view of the results, a larger number of items should be included to assess both the comfort and the organisational dimensions of hospital medical services.

The metric properties of the EMCA questionnaire, aimed at measuring satisfaction and perceived quality during hospitalisation in medical services, have shown to be valid and reliable for assessing these attributes, thus overcoming the methodological shortcomings of other instruments.⁶ The use of the EMCA questionnaire in other settings will enable external comparisons and benchmarking among hospitals.

The sample of cases analysed in this study, obtained from an acceptable response rate similar to that reported in the literature,^{23,24} came from a 4-year time series collected in nine public hospitals. The sample can therefore be considered representative, and ensures the generalisability of the results.

The overall value of 7.99 obtained for global satisfaction was high, and is similar to that reported in most of the literature, from classic studies such as the meta-analysis conducted by Hall and Dornan in 1998,²⁵ to the most recent research.^{24,26}

An analysis of the results for composite indicators of perceived quality revealed that the organisational and Information dimensions obtained the highest percentage of compliance satisfaction. In the case of the organisational dimension, this was due to high scores for aspects related to shift changes, continuity of care and management of visits, whilst for the dimension of information it was due to high scores for indicators related to clarity of information about the reason for admission, absence of contradictory information from staff, clear explanations of the treatment and care to be observed by the patient at home and the opportunity to ask doctors questions concerning the discharge report.

By contrast, the dimensions with the lowest percentage of compliance were comfort, mainly influenced by low scores for food and the general conditions of inpatient units, interpersonal manner and most especially the dimension of professional competence, which obtained the lowest level of compliance due to low scores for the professionalism of nursing and non-healthcare staff. However, a multivariate analysis revealed that these two dimensions, professional competence and interpersonal manner, together with Information, were the only ones which presented a statistically significant relationship with degree of satisfaction.

We coincided with most authors in considering information as one of the main determinants of satisfaction with inpatient care.²⁷⁻³⁰ The dimension of interpersonal manner, in particular as regards nursing staff, is also considered by several authors as an important determinant of satisfaction.^{23,27,30-32} As for organisational dimension, it is difficult to conduct a comparison of the results with respect to its influence on satisfaction, mainly due to the different components that are grouped together to form this dimension. With the exception of a few studies,^{2,30} and in contrast to the present study, this dimension is not usually considered an important predictor of satisfaction.

As in other studies, professional competence also appeared in this study as a determinant of satisfaction, although mainly in reference to the medical staff and almost always associated with the information received.^{27,28,30,33-35} However, other authors²³ have reported that the best predictor of satisfaction is treatment outcome, whereas Information has very little influence.

In this study a no statistically significant associations between sociodemographic variables and satisfaction level were found. These findings are similar to those obtained by authors such as Bjertnaes et al.,²⁶ and different from those reported in other studies^{23,27,36} where statistically significant associations were found between satisfaction and age, gender and/or educational level.

In this paper we did not find any association between previous hospital admissions to the same centre and satisfaction, although some authors such as Quintana²⁷ have reported that patients who have been admitted on earlier occasions tend to show a more critical attitude and present lower levels of satisfaction. It is necessary consider that satisfaction is a variable that usually shows a ceiling effect, that should be considered in future studies analysing the degree of insatisfacción.

A potential limitation of this study was not to obtain patients' opinions on the outcome of their hospitalisation, which has prevented an assessment of its influence on satisfaction with healthcare. It would be interesting to include items that inquire about reason for admission and outcome.

To conduct a metric analysis of the questionnaire, only those respondents who had completed all items were selected. Consequently, information corresponding to patients with missing data was not included in the multivariate analysis, which could suggest the existence of a possible selection bias in the results obtained. However, the lack of statistically significant differences with respect gender and the small effect size for age differences between the initial sample and the studied sample, indicate that the existence of such a bias can be virtually ruled out.

Another possible limitation of the study was that no information was collected regarding participants' prior health status, a variable that other studies have shown can influence the degree of satisfaction expressed.^{1,37}

In conclusion, the metric properties of the questionnaire used have been updated. The reliability values and percentage of explained variance were acceptable for all the dimensions analysed.

A valid and reliable questionnaire for assessing satisfaction in quality management programmes, both for internal purposes and for conducting external comparisons was obtained. This questionnaire is useful for clinicians and managers, enabling them to prioritise areas for intervention, those dimensions with greater room for improvement or those which have a greater impact on satisfaction.

A positive relationship between the degree of patient satisfaction and level of professional competence, interpersonal manner of healthcare staff and information received was found.

Conflict of interest

The authors declare no conflict of interest.

References

1. Crow R, Gage H, Hampson S, Hart J, Kimber A, Storey L, et al. The measurement of satisfaction with healthcare: implications for practice from a systematic review of the literature. *Health Technol Assess.* 2002;6.
2. Mira JJ, Rodríguez-Marín J, Peset R, Ybarra J, Pérez-Jover V, Palazón I, et al. Causas de satisfacción y de insatisfacción de los pacientes en hospitales y atención primaria. *Rev Calid Asist.* 2002;17:273-83.
3. Carminal J. La medida de la satisfacción: un instrumento de participación de la población en la mejora de la calidad de los servicios sanitarios. *Rev Calid Asist.* 2001;16:276-9.
4. Williams B. Patient satisfaction: a valid concept. *Soc Sci Med.* 1994;38:509-16.
5. Yogesh PP, Satyanarayana TC. Dimensions of hospital service quality: a critical review. *Int J Health Care Qual Assur.* 2013;26:308-40.
6. Serrano R, Loriente N. La anatomía de la satisfacción del paciente. *Salud Publica Mex.* 2008;50:162-72.
7. Yellen E, Galis G, Ricard R. The measurement of patient satisfaction. *J Nurs Care Qual.* 2002;16:23-9.
8. Bermejo RM, Hidalgo MD, Parra P, Más A, Gomis R. Propiedades métricas de las escalas de competencia profesional y trato del personal sanitario de servicios de consultas externas hospitalarias. *Psicothema.* 2011;23:295-300.
9. Contreras CW, Galarza CA. Perfil del paciente de medicina en un hospital público. *Acta Med Per.* 2009;26:212-6.
10. Fernández I, González E. Características clínicas y terapéuticas de los pacientes ingresados en un Servicio de Medicina Interna por insuficiencia cardíaca. *An Med Interna.* 2006;23:459-64.
11. Becchi MA, Belleli F, Clarizio M, Carulli N. Characteristics of patients admitted to a Division of Internal Medicine. Is hospital the best setting of care. *Ann Ital Med Int.* 2005;20:233-44.
12. San Román CM, Torremocha O, Medina P, Nuño E, Guil M, Barceló JM, et al. Estudio de pacientes pluripatológicos atendidos en el hospital comarcal de la Axarquía. *Rev Clin Esp.* 2001;201, extra 3:14.

13. Rodríguez J, Ruiz A, Gómez E, Ollero M, San Román C, Pujol E. El paciente pluripatológico en los servicios de Medicina Interna. *Rev Clin Esp.* 2001;201, extra 3:19.
14. García-Morillo García JS, Bernabeu M, Ollero M, Aguilar M, Ramírez N, González de la Puente MA, et al. Incidencia y características clínicas de los pacientes con pluripatología ingresados en una Unidad de Medicina Interna. *Med Clin.* 2005;25:5–9.
15. Zambrana JL, Velasco MJ, Díez F, Cruz G, Martín MD, Adarraga MD. Características clínicas diferenciales de los enfermos pluripatológicos hospitalizados en los servicios de Medicina interna. *Rev Clin Esp.* 2005;205:413–7.
16. Saturno PJ. Nueva encuesta posthospitalización del INSALUD. Murcia: Universidad de Murcia; 2001.
17. Reeves R, Coulter A, Jenkinson C, Cartwright J, Bruster S, Richards N. Development and pilot testing of questionnaires for use in the acute NHS inpatient survey programme. Report Picker Institute Europe; 2002.
18. Aday LA, Cornelius LJ. Designing and conducting health surveys: a comprehensive guide. 3rd ed. San Francisco: Josey-Bass Pub.; 2006.
19. Cohen J. Statistical power analysis for the behavioral sciences. Hillsdale, NJ: Lawrence Erlbaum Associates; 1998.
20. Landrum M, Bronskill S, Normand S. Analytic methods for constructing cross-sectional profiles of health care providers. *Health Serv Res Methodol.* 2000;200:23–48.
21. SPSS. Statistical package for social sciences for windows, version 19.0. Chicago IL: SPSS Inc.; 2006.
22. Granado S, Rodríguez C, Olmedo MC, Chacón A, Vigil D, Rodríguez P. Diseño y validación de un cuestionario para evaluar la satisfacción de los pacientes atendidos en las consultas externas de un hospital de Madrid en 2006. *Rev Esp Salud Publica.* 2007;81:637–45.
23. Schoenfelder T, Klewer J, Kugler J. Determinants of patient satisfaction: a study among 39 hospitals in an in-patient setting in Germany. *Int J Qual Health Care.* 2011;23:503–9.
24. Elliot M, Lehrman WG, Beckett M, Goldstein E, Hambarsoomina K, Giordano. Gender differences in patients' perceptions of inpatient care. *Health Serv Res.* 2012;47:1482–501.
25. Hall J, Dornan M. Meta-analysis of satisfaction with medical care: description of research domain and analysis of overall satisfaction levels. *Soc Sci Med.* 1988;27:637–44.
26. Bjertnaes O, Strømseng I, Hestad H. Overall patient satisfaction with hospitals: effects of patient-reported experiences and fulfilment of expectations. *BMJ Qual Saf.* 2012;21:39–46.
27. Quintana JM, Predáez N, Bilbao A, Aizpuru F, Escobar A, Esteban C, et al. Predictors of patient satisfaction with hospital health care. *BMC Health Serv Res.* 2006;6:102.
28. González N, Quintana JM, Bilbao A, Escobar A, Aizpuru F, Thompson A, et al. Development and validation of an inpatient satisfaction questionnaire. *Int J Qual Health Care.* 2005;17:465–72.
29. Dierssen T, Rodríguez P, Robles M, Brugos V, Gómez I, Llorca J. Factores que determinan la alta satisfacción del usuario con la asistencia hospitalaria. *An Sist Sanit Navar.* 2009;32:317–25.
30. Castillo L, Dougna A, Vicente I, Muñoz V, Rojas V. Los predictores de satisfacción de pacientes en un centro hospitalario universitario. *Rev Med Chil.* 2007;135:696–701.
31. Otani K, Herrmann P, Kurz R. Improving patients satisfaction in hospital care setting. *Health Serv Manage Res.* 2011;24:163–9.
32. Ortiz RM, Muñoz S, Torres E. Satisfacción de los usuarios de 15 hospitales de Hidalgo, México. *Rev Esp Salud Publica.* 2004;78:527–37.
33. Parra P, Bermejo RM, Más A, Hidalgo MD, Gomis R, Calle JE. Factores relacionados con la satisfacción del paciente en los servicios de urgencias hospitalarios. *Gac Sanit.* 2012;26:159–65.
34. Murillo C, Saurina C. Medida de la importancia de las dimensiones de la satisfacción en la provisión de servicios de salud. *Gac Sanit.* 2013;27:304–9.
35. Morales JM, Bonill de las Nieves C, Celdrán M, Morilla JC, Martín FJ, Contreras E. Diseño y validación de instrumento de evaluación de la satisfacción con los servicios de atención domiciliaria: SATISFAD. *Gac Sanit.* 2007;21:106–13.
36. González N, Quintana JM, Bilbao A, Esteban C, San Sebastian JA, de la Sierra E, et al. Satisfacción de los usuarios de 4 hospitales del Servicio Vasco de Salud. *Gac Sanit.* 2008;22:210–7.
37. Xiao H, Barber JP. The effect of perceived health status on patient satisfaction. *Value Health.* 2008;11:719–25.